

Accelerated telomere shortening in response to life stress

Elissa S. Epel^{*†}, Elizabeth H. Blackburn[‡], Jue Lin[‡], Firdaus S. Dhabhar[§], Nancy E. Adler^{*}, Jason D. Morrow[¶], and Richard M. Cawthon^{||}

^{*}Department of Psychiatry, University of California, 3333 California Street, Suite 465, San Francisco, CA 94143; [‡]Department of Biochemistry and Biophysics, University of California, San Francisco, CA 94143; [§]Department of Oral Biology, College of Dentistry, and Department of Molecular Virology, Immunology, and Medical Genetics, College of Medicine, Ohio State University, Columbus, OH 43210; [¶]Department of Medicine and Pharmacology, Vanderbilt University School of Medicine, Nashville, TN 37232; and ^{||}Department of Human Genetics, University of Utah, 15 North 2030 E Street, Room 2100, Salt Lake City, UT 84112

Contributed by Elizabeth H. Blackburn, September 28, 2004

Numerous studies demonstrate links between chronic stress and indices of poor health, including risk factors for cardiovascular disease and poorer immune function. Nevertheless, the exact mechanisms of how stress gets “under the skin” remain elusive. We investigated the hypothesis that stress impacts health by modulating the rate of cellular aging. Here we provide evidence that psychological stress—both perceived stress and chronicity of stress—is significantly associated with higher oxidative stress, lower telomerase activity, and shorter telomere length, which are known determinants of cell senescence and longevity, in peripheral blood mononuclear cells from healthy premenopausal women. Women with the highest levels of perceived stress have telomeres shorter on average by the equivalent of at least one decade of additional aging compared to low stress women. These findings have implications for understanding how, at the cellular level, stress may promote earlier onset of age-related diseases.

psychological stress | telomere length | telomerase | oxidative stress

People who are stressed over long periods tend to look haggard, and it is commonly thought that psychological stress leads to premature aging and the earlier onset of diseases of aging. Numerous studies demonstrate links between chronic stress and indices of poor health, including risk factors for cardiovascular disease and poorer immune function (1, 2). Nevertheless, the exact mechanisms of how such stress exerts these effects are not well known, including whether stress accelerates aging at a cellular level and how cellular aging translates to organismal aging. Recent research points to the crucial roles of telomeres and telomerase in cellular aging and potentially in disease. Telomeres are DNA-protein complexes that cap chromosomal ends, promoting chromosomal stability. When cells divide, the telomere is not fully replicated because of limitations of the DNA polymerases in completing the replication of the ends of the linear molecules, leading to telomere shortening with every replication (3). *In vitro*, when telomeres shorten sufficiently, the cell is arrested into senescence. In people, telomeres shorten with age in all replicating somatic cells that have been examined, including fibroblasts and leukocytes (4). Thus, telomere length can serve as a biomarker of a cell's biological (versus chronological) “age” or potential for further cell division.

Telomerase, a cellular enzyme, adds the necessary telomeric DNA (T₂AG₃ repeats) onto the 3' ends of the telomere (5). Telomerase also has direct telomere-protective functions (6). In human T cells, telomerase activity increases with acute antigen exposure but decreases with repeated antigen stimulation and as the cells approach senescence (7). People with dyskeratosis congenita, a rare genetic disease that diminishes the ability to synthesize sufficient telomerase, have shortened telomeres and die prematurely from progressive bone marrow failure and vulnerability to infections (8).

Cellular environment also plays an important role in regulating telomere length and telomerase activity. Most notably, *in vitro*, oxidative stress can shorten telomeres and antioxidants can decelerate shortening (9, 10). Perceived stress has been linked to one measure of oxidative DNA damage in leukocytes in women (11, 12). Given these observed links, we hypothesized that chronic psychological stress may lead to telomere shortening and lowered telomerase function in peripheral blood mononuclear cells (PBMCs) and to oxidative stress.

Methods

To study both objective (event-/environment-based) and subjective (perception-based) stress, we examined 58 healthy premenopausal women who were biological mothers of either a healthy child ($n = 19$, “control mothers”) or a chronically ill child ($n = 39$, “caregiving mothers”). The latter were predicted to have, on average, greater environmental exposure to stress. Women in both groups completed a standardized 10-item questionnaire assessing level of perceived stress over the past month (13). This design allowed us to examine the importance of perceived stress and measures of objective stress (caregiving status and chronicity of caregiving stress based on the number of years since a child's diagnosis) (see *Supporting Text*, which is published as supporting information on the PNAS web site). All analyses were conducted controlling for age because we wanted to test for telomere shortening caused by stress independent of the women's chronological age (age was related to telomere length, $r = -0.23$, $P < 0.04$).

Each subject was 20–50 years old [mean (M) = 38 ± 6.5 years] and had at least one biological child living with her. Subjects were free of any current or chronic illness (see *Supporting Text*). Use of oral contraceptives was similar in the caregiver and control groups. Obesity level was quantified by body-mass index (BMI): weight (in kilograms)/[height (in meters) × height (in meters)]. Blood was drawn in a fasting state on a morning during the first 7 days of the follicular stage of the menstrual cycle.

Mean telomere length and telomerase activity were measured quantitatively in the PBMCs that were stored frozen at -80°C . Telomere length values were measured from DNA by a quantitative PCR assay that determines the relative ratio of telomere repeat copy number to single-copy gene copy number (T/S ratio) in experimental samples as compared with a reference DNA sample (14). Telomerase activity was measured by the telomerase repeat amplification protocol (15) with a commercial kit (Trapeze, Chemicon) and all values used were in the linear quantitative range. Vitamin E (alpha-tocopherol) was measured

Freely available online through the PNAS open access option.

Abbreviations: PBMC, peripheral blood mononuclear cell; BMI, body-mass index; T/S, telomere repeat copy number/single-copy gene copy number.

[†]To whom correspondence should be addressed. E-mail: elissa@itsa.ucsf.edu.

© 2004 by The National Academy of Sciences of the USA




Fig. 2. Telomere length and telomerase activity levels in extreme high- and low-stress groups. (A) Average telomere length and SE are shown. The high-stress group had shorter telomeres even after controlling for age and BMI [$F(1, 27) = 12.8, P < 0.001$]. (B) Average telomerase activity and SE are shown. The high-stress group had lower telomerase activity even after controlling for age and BMI [$F(1, 27) = 3.1, P < 0.05$]. Because values for telomerase activity were not normally distributed, the values were converted to follow a normal distribution by using a natural logarithm transformation. All statistical analyses were performed on the transformed variable. The raw (unadjusted) levels of telomerase are presented throughout because these values are more meaningful than transformed values.

age and BMI. Independent of their chronological age and BMI, the high-stress group had significantly shorter telomeres (raw mean T/S ratio = 1.13 ± 0.17), which converts to 3,110-bp telomeres than the low-stress group (raw mean T/S ratio = 1.33 ± 0.15), which converts to 3,660-bp telomeres (Fig. 2A).

Because telomere length declines during normal aging, one can estimate the years of aging expected to pass to bring the relatively long telomeres of the bottom-stress-quartile individuals down to the relatively short lengths of the top-stress-quartile individuals. Cross-sectional studies of age versus telomere length in the general population show that telomere shortening in young adulthood appears minimal in lymphocytes or PBMCs (4, 17–19) and becomes faster (≈ 60 bp/year) in older age (50–70 years) (17, 19). There are no studies that estimate average expected telomere shortening in the exact age range studied here (20–50 years), which largely falls within the young adulthood period of minimal loss. Therefore, in translating telomere shortening to years of aging, we based our estimates on studies averaging telomere shortening across adulthood (ages 20–95 years), which find a loss of 31–63 bp/year (17, 20, 21). Thus, the 550-bp shortening in the high-stress group indicates that their lymphocytes had aged the equivalent of 9–17 additional years, compared with the low-stress group.

Strikingly, the high-stress group also had significantly lower telomerase activity (Fig. 2B) and higher oxidative stress than the low-stress group. The mean telomerase activity, adjusted for BMI and age, was 48% lower in the high-stress group ($M = 0.053 \pm 0.016$ versus 0.092 ± 0.016 telomerase units per 10,000 cells; $F(1, 27) = 3.1; P < 0.045$). The mean oxidative stress index was $F(3, 22) = 4.5, P < 0.025$ ($M = 0.052 \pm 0.010$ versus 0.085 ± 0.010 arbitrary units of isoprostanes/vitamin E). When the results were analyzed by high versus low telomere length, the conclusions were similar in that the short telomere length group had higher perceived stress, higher oxidative stress, and lower telomerase activity (see *Supporting Text*).

Discussion

The exact mechanisms that connect the mind to the cell are unknown. Although it is well accepted that cell senescence can include stress-induced processes, psychological stress has not yet been considered as part of the stress pathway. The current findings suggest that stress-induced premature senescence in people might be influenced by chronic or perceived life stress. Psychological stress could affect cell aging through at least three nonmutually exclusive pathways: immune cell function or distribution, oxidative stress, or telomerase activity. We considered whether stress might have decreased naïve T cells and increased memory T cells [which have shorter telomere length (22)], but the data did not support this (Table 2, which is published as supporting information on the PNAS web site). Second, stress could potentially lead to oxidative stress by means of chronic activation of the autonomic and neuroendocrine stress responses. Although this hypothesis has never been tested *in vivo*, the relationship between stress hormones and oxidative stress has been clearly demonstrated at the cellular level. Glucocorticoids, the primary adrenal hormones secreted during stress, increase oxidative stress damage to neurons, in part by increasing glutamate and calcium and decreasing antioxidant enzymes (23, 24). It is also notable that, in women, self-reported distress has been related to greater oxidative DNA damage (8-OH-dG) (12). Oxidative stress shortens telomeres in cells cultured *in vitro* (10). Our findings that perceived and chronic stress correlated with higher oxidative stress and shorter telomere length demonstrate this relationship cross-sectionally for the first time *in vivo*. Lastly, if the observed lowered telomerase activity represents chronic levels, it too could have contributed to the shortened telomeres in PBMCs.

Although being a caregiver *per se* was not related to telomere length, the chronicity of caregiving stress was related to telomere length. It is a formal possibility that the association between stress and telomere length reported here is in part due to people with longer telomeres being psychologically more resistant to objective stressors than people with shorter telomeres. Engineering worms to have longer telomeres resulted in an extension of their life spans, even though all of their somatic cells are postmitotic and their telomeres do not shorten during aging (25). Further, the biochemical pathway involved in this effect of telomere length on life span is the same pathway previously shown to mediate increased physiological stress resistance in response to various genetic variants that increase life span. It is unknown whether physiological stress resistance is related to psychological stress resistance. Perceived stress may be either causally or correlationally related to telomere length. However, the relationship between chronicity of caregiving stress (an objectively based measure) and telomere length implies that stress preceded telomere shortening, because telomere length cannot have influenced the number of years one has been a caregiver. Clearly, longitudinal studies in which telomere length is assayed repeatedly are needed to directly test whether the rate of telomere shortening in individuals with higher levels of reported stress is actually faster than in individuals with lower

stress. Further, clinical trials of interventions effective in lowering perceived stress in adult life could test whether they slow the rate of telomere shortening.

These associations between stress and cell aging have clinically significant implications for human health. A 50% deficiency in telomerase RNA gene dosage caused by the rare genetic disease dyskeratosis congenita is sufficient to cause premature death in adults from bone marrow failure and vulnerability to infections (8). In the elderly, telomere shortening is strongly associated with higher mortality rates (26). Lastly, patients with early myocardial infarction had leukocyte telomere lengths that were equivalent to those typical of a person \approx 11 years older than controls, similar to the magnitude of accelerated cell aging observed in our high-stress group (27).

In summary, in healthy women, psychological stress is associated with indicators of accelerated cellular and organismal aging: oxidative stress, telomere length, and telomerase activity in PBMCs. Many questions remain, such as whether shorter telomeres in leukocytes lead to earlier immune senescence and relate to shorter telomeres in other proliferative cells, such as cardiovascular endothelium. Nonetheless, although the exact pathway of events from perception of stress in the brain to

somatic cell longevity is unclear, the results reported here now implicate shorter telomeres in the adverse health *sequelae* of prolonged psychological stress.

We thank Drs. Teresa Seeman, Margaret Kemeny, and Bruce McEwen for intellectual support; Jean M. Tillie for expert help with flow cytometry; Michael Acree for statistical expertise; Drs. Melvin Heyman, Bryna Siegel, and Paul Harmatz for help recruiting clinic participants; Sheryln Jimenez, Denise Kruszewski, and Drs. Judith Moskowitz, Susan Folkman, and Judith Stewart for crucial help; and the busy mothers of volunteering their time. E.S.E. was supported by the John D. and Catherine T. MacArthur Foundation Network on Socioeconomic Status and Health; the Hellman Family Fund; and the University of California, San Francisco, Pediatric Clinical Research Center (under the auspices of National Institute of Mental Health Grant M01-RR01271), National Institute of Mental Health Award K08 MH64110-01A1, and a National Alliance for Research on Schizophrenia and Depression Young Investigator's Award. E.H.B. was supported by the Steven and Michele Kirsch Foundation and National Institutes of Health Grant GM26259. J.D.M. was supported by the Burroughs Wellcome Fund Clinical Scientist Award in Translational Research and National Institutes of Health Grants GM15431, CA77839, DK48851, and RR00095. F.S.D. was supported by the Dana Foundation and National Institutes of Health Grant AI48995.

- McEwen, B. (1998) *N. Engl. J. Med.* **338**, 171–179.
- Segerstrom, S. & Miller, G. (2004) *Psychol. Bull.* **130**.
- Chan, S. R. & Blackburn, E. H. (2004) *Philos. Trans. R. Soc. London B* **359**, 109–121.
- French, R., Blackburn, E. & Shannon, K. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 5607–5610.
- Blackburn, E. H. (1997) *Biochemistry (Moscow, Russ. Fed.)* **62**, 1196–1201.
- Chan, S. W. & Blackburn, E. H. (2003) *Mol. Cell* **11**, 1379–1387.
- Weng, N. P., Palmer, L. D., Levine, B. L., Lane, H. C., June, C. H. & Hodes, R. J. (1997) *Immunol. Rev.* **160**, 43–54.
- Vulliamy, T., Marrone, A., Goldman, F., Dearlove, A., Bessler, M., Mason, P. J. & Dokal, I. (2001) *Nature* **413**, 432–435.
- von Zglinicki, T., Saretzki, G., Docke, W. & Lotze, C. (1995) *Exp. Cell Res.* **220**, 186–193.
- von Zglinicki, T. (2002) *Trends Biochem. Sci.* **27**, 339–344.
- Irie, M., Asami, S., Nagata, S., Miyata, M. & Kasai, H. (2001) *Int. Arch. Occup. Environ. Health* **74**, 153–157.
- Irie, M., Asami, S., Ikeda, M. & Kasai, H. (2003) *Biochem. Biophys. Res. Commun.* **311**, 1014–1018.
- Cohen, S. & Williamson, G. (1988) in *The Social Psychology of Health: Claremont Symposium on Applied Social Psychology*, ed. Oskamp, S. S. S. (Sage, Newbury Park, CA).
- Cawthon, R. M. (2002) *Nucleic Acids Res.* **30**, e47.
- Kim, N. W. & Wu, F. (1997) *Nucleic Acids Res.* **25**, 2595–2597.
- Morrow, J. D. & Roberts, L. J., II (2002) *Methods Mol. Biol.* **186**, 57–66.
- Iwama, H., Ohyashiki, K., Ohyashiki, J. H., Hayashi, S., Yahata, N., Ando, K., Toyama, K., Hoshika, A., Takasaki, M., Mori, M. & Shay, J. W. (1998) *Hum. Genet.* **102**, 397–402.
- Jeanclous, E., Schork, N. J., Kyvik, K. O., Kimura, M., Skurnick, J. H. & Aviv, A. (2000) *Hypertension* **36**, 195–200.
- Rufer, N., Brummendorf, T. H., Kolvraa, S., Bischoff, C., Christensen, K., Wadsworth, L., Schulzer, M. & Lansdorp, P. M. (1999) *J. Exp. Med.* **190**, 157–167.
- Hastie, N. D., Dempster, M., Dunlop, M. G., Thompson, A. M., Green, D. K. & Allshire, R. C. (1990) *Nature* **346**, 866–868.
- Satoh, H., Hiyama, K., Takeda, M., Awaya, Y., Watanabe, K., Ihara, Y., Maeda, H., Ishioka, S. & Yamakido, M. (1996) *Jpn. J. Hum. Genet.* **41**, 413–417.
- Hodes, R. J., Hathcock, K. S. & Weng, N. P. (2002) *Nat. Rev. Immunol.* **2**, 699–706.
- McIntosh, L. J. & Sapolsky, R. M. (1996) *Neurotoxicology* **17**, 873–882.
- Patel, R., McIntosh, L., McLaughlin, J., Brooke, S., Nimon, V. & Sapolsky, R. (2002) *J. Neurochem.* **82**, 118–125.
- Joeng, K. S., Song, E. J., Lee, K. J. & Lee, J. (2004) *Nat. Genet.* **36**, 607–611.
- Cawthon, R., Smith, K., O'Brien, E., Sivatchenko, A. & Kerber, R. (2003) *Lancet* **361**, 393–395.
- Brouillette, S., Singh, R. K., Thompson, J. R., Goodall, A. H. & Samani, N. J. (2003) *Arterioscler. Thromb. Vasc. Biol.* **23**, 842–846.